



# Priconazole<sup>®</sup> Lotion & Spray, 1% (miconazole nitrate)



- FDA approved, ANADA 200-362
- Priconazole<sup>®</sup> Lotion & Spray, 1% are approved by the FDA as equivalent to the pioneer product Conofite<sup>®</sup> Lotion 1%.<sup>1</sup>
- Priconazole<sup>®</sup> Lotion & Spray, 1% are 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight).
- Priconazole<sup>®</sup> Lotion, 1% is supplied in 60 mL (2 fl oz) plastic bottles with dropper plugs and Priconazole<sup>®</sup> Spray, 1% is supplied in 120 mL (4 fl oz) and 240 mL (8 fl oz) plastic bottles with sprayers.

A synthetic antifungal agent for use on dogs and cats. Indicated for treatment of *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*. Please consult package insert for full product information (see back page).

Size	Reorder No.	Lbs/Case	Case Pack
60 mL (2 oz)	TP321	13 lbs.	72 x 1
120 mL (4 oz)	TP322	4 lbs.	12 x 1
240 mL (8 oz)	TP323	7 lbs.	12 x 1

NDC# 58829-297-60

**INDICATIONS:** For the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

**DOSAGE:** Apply a light covering to affected areas, once daily for 2 to 4 weeks. See accompanying literature for full directions.

Avoid contact with eyes, since irritation may result.

Wash hands thoroughly after administration to avoid spread of fungal infection.

TAKE TIME OBSERVE LABEL DIRECTIONS

Priority Care and Priconazole are registered trademarks of First Priority, Inc.  
Rev. 04-08

## Priconazole<sup>®</sup> Lotion, 1%

(miconazole nitrate)

Topical antifungal agent for dogs and cats.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**For Animal Use Only • Keep Out of Reach of Children**

Net Contents: 60 mL (2 fl oz)

ANADA 200-362, Approved by FDA

Manufactured By:  
FIRST PRIORITY, INC., Elgin, IL 60123-1146  
www.prioritycare.com

**ACTIVE INGREDIENT:**  
Miconazole Nitrate.....1.15%  
(equivalent to 1% miconazole base by weight)

**STORAGE:** Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.

Lot No.      Exp. Date

Reorder No. TP321PC

NOT RETURNABLE IF OUTDATED  
  
 6230812976016

<sup>1</sup> Conofite<sup>®</sup> is a registered trademark of Johnson & Johnson Corporation, New Brunswick, NJ

For a complete list of Priority Care distributors, please call our Customer Service Department at: 1-800-650-4899, or visit our Website: [www.prioritycare.com](http://www.prioritycare.com)

# Miconazole Nitrate Lotion, 1% & Spray, 1%

(miconazole nitrate)

Topical antifungal agent for dogs and cats.

**DESCRIPTION:** Miconazole Nitrate Lotion, 1% & Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

**INDICATIONS:** Miconazole Nitrate Lotion, 1% & Spray, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

**PRECAUTIONS:** In the event of sensitization or irritation due to Miconazole Nitrate Lotion, 1% & Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

**DOSAGE AND ADMINISTRATION:** Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

**Lotion:** Apply a light covering of Miconazole Nitrate Lotion, 1% (miconazole nitrate) to affected areas, once daily, for 2 to 4 weeks. Application is best accomplished using a gauze pad or cotton swab.

**Spray:** Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

**HOW SUPPLIED:** Miconazole Nitrate Lotion, 1% (miconazole nitrate) is available in 30 mL & 60 mL bottles with droppers, and Miconazole Nitrate Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Iss. 04-09

Manufactured by:  
First Priority, Inc.  
Elgin, IL U.S.A.  
[www.prioritycare.com](http://www.prioritycare.com)



ANADA# 200-362,  
Approved by FDA