



Privermectin® Drench for Sheep

(ivermectin)



- FDA approved, ANADA 200-327
- Privermectin® Drench for Sheep is approved by the FDA as equivalent to the pioneer product Ivermec® Drench for Sheep.¹
- Privermectin® Drench for Sheep is supplied in 240 mL (8 fl oz) and 960 mL (32.46 fl oz) plastic bottles. The 960 mL bottle will easily facilitate the use of current dispensing instruments.
- 83 - 100 lb. doses.

A ready-to-use, free-flowing solution of ivermectin for the treatment and control of worms and bots in sheep. Please consult package insert for full product information (see back page).

<u>Size</u>	<u>Reorder No.</u>	<u>Lbs/Case</u>	<u>Case Pack</u>
240 mL	OM041	5 lbs.	6 x 1
960 mL	OM040	11 lbs.	4 x 1



¹ Ivermec® is a registered trademark of Merial LLC, Essex, United Kingdom

For a complete list of Priority Care distributors, please call our Customer Service Department at: 1-800-650-4899, or visit our Website: www.prioritycare.com

Privermectin® Drench for Sheep
0.08% Solution
 (ivermectin)

Approved by FDA under ANADA # 200-327

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE.

PRODUCT DESCRIPTION: Privermectin® Drench for Sheep is a ready-to-use, free-flowing solution of ivermectin. It is formulated to deliver the recommended dose rate of 0.2 mg ivermectin per 1 kg body weight given orally at a volume of 3.0 mL per 26 lbs body weight.

INDICATIONS: Privermectin® Drench for Sheep provides treatment and control of adult and fourth-stage larvae of the following parasites: Gastrointestinal Roundworms – *Haemonchus contortus*, *Ostertagia circumcincta*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia curtipice*, *Nematodirus spathiger*, *N. battus*, and *Oesophagostomum columbianum*; Lungworms – *Dictyoacaulus filaria*; and all the larval stages of Nasal Bot – *Oestrus ovis*. It also provides treatment and control of adult forms only of the following Gastrointestinal Roundworms – *Haemonchus placei*, *Cooperia oncophora*, *Strongyloides papillosus*, *Oesophagostomum venulosum*, *Trichouris ovis*, and *Chabertia ovis*.

1

or ground water. Do not contaminate water by direct application or by the improper disposal of drug containers. Spills should be contained and soaked up with absorbent towels or into loose soil. Gloves should be worn to prevent skin exposure. All the collected materials (contaminated towels and soil), as well as all empty drug containers, should be placed in an impervious film (plastic) bag and disposed of by incineration or in an approved landfill.

STORAGE INFORMATION: Store at 68° - 77° F (20° - 25° C). Excursions between 59° - 86° F (15° - 30° C) are permitted.

Approved by FDA under # ANADA 200-327

Rev. 02-20

Reorder No. 0M040PC

4

DOSAGE AND ADMINISTRATION: Privermectin® Drench for Sheep may be used in any standard drenching equipment or in any equipment which provides a consistent dose volume. Privermectin® Drench for Sheep is administered orally at a dose of 3.0 mL (2.4 mg ivermectin) per 26 lbs body weight or 200 mcg ivermectin per kilogram of body weight.

Coughing may be observed in some animals during and for several minutes following drenching.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

RESIDUE WARNING: Do not treat sheep within 11 days of slaughter.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse effects in users, to obtain an SDS, or for assistance call 1-800-650-4899.

PRECAUTIONS: Privermectin® (ivermectin) Drench for Sheep has been formulated for use in sheep only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

OTHER WARNINGS: Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers.

Treatment with a dewormer used in conjunction with parasite management

2

practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance.

Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd/flock, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method).

A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

Keep this and all drugs out of reach of children.


Refrain from smoking and eating when handling. Avoid contact with eyes. Immediately wash hands and any spills on the skin with plenty of soap and water following use.

Restricted Drug (California) - Use Only as Directed.

Environmental Safety: Studies indicate that when ivermectin comes in contact with the soil, it readily and tightly binds to the soil and becomes inactive over time. Free ivermectin may adversely affect fish and certain water-born organisms on which they feed. Do not permit water runoff from feedlots to enter lakes, streams,

3

NDC# 58829-293-32



Privermectin® Drench for Sheep


(ivermectin)

0.08% Solution
Parasiticide

For the Treatment and Control
of Worms and Bots of Sheep

- 83-100 lb Doses •

See booklet for additional information.



Rev. 02-20


For Animal Use Only • Keep Out of Reach of Children

Net Contents: 32.46 fl oz (1 qt 0.46 fl oz) 960 mL


Approved by FDA under ANADA # 200-327

Manufactured By
FIRST PRIORITY, INC., Elgin, IL 60123

Made in U.S.A. www.prioritycare.com



Priority Care and Privermectin are registered trademarks of First Priority, Inc.



TAKE TIME

5