





Priconazole[®] Lotion & Spray, 1%

(miconazole nitrate)






- Approved by FDA under <u>ANADA # 200-362</u>
- Priconazole[®] Lotion & Spray, 1% are 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight).
- Priconazole[®] Lotion, 1% is supplied in 60 mL (2 fl oz) plastic bottles with dropper plugs and Priconazole[®] Spray, 1% is supplied in 120 mL (4 fl oz) and 240 mL (8 fl oz) plastic bottles with sprayers.

A synthetic antifungal agent for use on dogs and cats. Indicated for treatment of Microsporum canis, Microsporum gypseum, and Tricophyton mentagrophytes. Please consult package insert for full product information (see back page).

<u>Size</u>	Reorder No.	Lbs/Case	<u>Case Pack</u>
60 mL (2 oz)	TP321	13 lbs.	72 x 1
120 mL (4 oz)	TP322	4 lbs.	12 x 1
240 mL (8 oz)	TP323	7 lbs.	12 x 1



For a complete list of Priority Care distributors, please call our Customer Service Department at: 1-800-650-4899, or visit our Website: www.prioritycare.com

Priconazole[®] Spray, 1% (miconazole nitrate)

Topical antifungal agent for dogs and cats.

DESCRIPTION: Priconazole[®] Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

INDICATIONS: Priconazole[®] Spray, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

PRECAUTIONS: In the event of sensitization or irritation due to Priconazole[®] Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

DOSAGE AND ADMINISTRATION: Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

Spray: Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

HOW SUPPLIED: Priconazole[®] Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

CONTACT INFORMATION: To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc at (800) 650-4899. Additional information can be found at www.prioritycare.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae

Rev. 08-22

Manufactured by: First Priority, Inc. Elgin, IL 60123 www.prioritycare.com Approved by FDA under ANADA # 200-362



Priority Care and Priconazole are registered trademarks of First Priority, Inc.