





Privermectin[®] Drench for Sheep

(ivermectin)



- Approved by FDA under <u>ANADA # 200-327</u>
- Privermectin[®] Drench for Sheep is approved by the FDA as equivalent to the pioneer product Ivomec[®] Drench for Sheep.¹
- Privermectin[®] Drench for Sheep is supplied in 240 mL (8 fl oz) and 960 mL (32.46 fl oz) plastic bottles. The 960 mL bottle will easily facilitate the use of current dispensing instruments.
- 83 100 lb. doses.

A ready-to-use, free-flowing solution of ivermectin for the treatment and control of worms and bots in sheep. Please consult package insert for full product information (see back page).

<u>Size</u>	<u>Reorder No</u>	Lbs/Case	<u>Case Pack</u>
240 mL	OM041	5 lbs.	6 x 1
960 mL	OM040	11 lbs.	4 x 1



¹Ivomec is a registerted trademark of Boehringer Ingelheim Animal Health USA Inc.

For a complete list of Priority Care distributors, please call our Customer Service Department at: 1-800-650-4899, or visit our Website: www.prioritycare.com

 practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd/flock, prior to the use of any dewormer, effectiveness of and indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring. Keep this and all drugs out of reach of children. Restricted Drug out of reach of children. Restricted Drug (California) - Use Only as Directed. Restricted Drug (California) - Use Only as Directed. Restricted Drug (California) - Use Only as Directed. Fer ivermectin may adversely affect fish and certain water-born organisms on which they feed. Do not permit water runoff from feedlots to enter lakes, streams, which they feed. Do not permit water runoff from feedlots to enter lakes, streams, which they feed. Do not permit water runoff from feedlots to enter lakes, streams, enterpermited advected in the sole. 	<image/> <image/> <section-header><section-header><section-header><section-header><text><text><text></text></text></text></section-header></section-header></section-header></section-header>
 DOSAGE AND ADMINISTRATION: Privermectin[®] Drench for Sheep may be used in any standard drenching equipment or in any equipment which provides a consistent dose volume. Privermectin[®] Drench for Sheep is administered orally at a dose of 3.0 mL (2.4 mg ivermectin) per 26 lbs body weight or 200 mcg ivermectin per kilogram of body weight. Coughing may be observed in some animals during and for several minutes following drenching. Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance. The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To eacort adverse effects in users, to obtain an SDS, or for assistance call to To eacort adverse effects in users, to obtain an SDS, or for assistance call to To report adverse effects in users, to obtain an SDS, or for assistance call to To eacort adverse effects in users, to obtain an SDS, or for assistance call to Eaco-650-4899 	Priority Care and Privermectin are The Mark of First Priority, Inc.
 Privermectin[®] Drench for Sheep 0.08% Solution (ivermectin) Approved by FDA under ANADA # 200-327 (ivermectin) Approved by FDA under ANADA # 200-327 Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. COR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ORAL USE IN SHEEP ONLY. NOT FOR HUMAN USE. FOR ONLY OF EVENTIATION: Privermectin[®] Drench for Sheep is a ready-to-use, if a columption of ivermectin. FOR ONLY DESCRIPTION: Privermectin[®] Drench for Sheep provides treatment and control of adult and fourth-stage larvae of the following parasites: Gastrointestinal Roundworms - Haemonchus contortus, Ostertagia circumcincta, Trichostrongylus axei, T. colubriformis, Cooperia curticei, Nernatodirus spathiger, M. battus, and Oesopha- gostommum columbianum. Lungworms - Dichyocaulus finaria; and all the larval stages of Masel Bot - Oestrus ovis, it also provides treatment and control of adult forms only of the following Gastrointestinal Roundworms - Haemonchus placei, Cooperia oncophora, Strongyloides papillosus, Oesophagostomum venulosum, Trichuris ovis, and Chabertia ovina. 	or ground water. Do not contaminate water by direct application or by the improper disposal of drug containers. Spills should be contained and soaked up with absorbent towels or into loose soil. Gloves should be worn to prevent skin exposure. All the collected materials (contaminated towels and soil), as well as all empty drug containers, should be placed in an impervious film (plastic) bag and disposed of by incineration or in an approved landfill. StoRadE INFORMATION: Store at 68° - 77° F (20° - 25° C). Excursions between 59° - 86° F (15° - 30° C) are permitted. Approved by FDA under # ANADA 200-327 Rev. 03-22 Rev. 03-22 Reorder No. 0M040PC