




Sulfadimethoxine Soluble Powder

Antibacterial Soluble Powder


R_x



An antibacterial soluble powder for use in broiler and replacement chickens, meat-producing turkeys, and dairy calves, dairy heifers, and beef cattle.

For the treatment of disease outbreaks of coccidiosis, fowl cholera and infectious coryza in broiler and replacement chickens, for disease outbreaks of coccidiosis and fowl cholera in meat-producing turkeys, and for treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Sphaerophorus necrophorus* sensitive to sulfadimethoxine in dairy calves, dairy heifers, and beef cattle

<u>Size</u>	<u>Reorder No</u>	<u>Lbs/Case</u>	<u>Case Pack</u>
107 g	OM300	6.5 lbs.	25 x 1 pail

Priority Care  NDC# 58829-348-03

Sulfadimethoxine Soluble Powder

Antibacterial Soluble Powder

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each packet contains 3.34 oz (94.6 g) sulfadimethoxine in the form of the soluble sodium salt and disodium edetate.

Restricted Drug (California) - Use Only as Directed. Not For Human Use.

For Broiler and Replacement Chickens Only - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

For Meat-producing Turkeys Only - Use for the treatment of disease outbreaks of coccidiosis and fowl cholera.

For Dairy Calves, Dairy Heifers, and Beef Cattle - Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Sphaerophorus necrophorus* sensitive to sulfadimethoxine.

Not for Use in Humans • Keep Out of Reach of Children

Iss. 11-24
Reorder No. OM300PC
Net Contents: 3.77 oz (107 g)
Approved by FDA under ANADA # 200-443

NOT RETURNABLE IF OUTDATED

Made in U.S.A. Manufactured By: FIRST PRIORITY, INC., Elgin, IL www.prioritycare.com

Sulfadimethoxine Soluble Powder

Dosage and Administration:

Species	Concentration	Use Directions
Chickens	0.05%	Contents of packet to 50 gallons of water.
Turkeys	0.025%	Contents of packet to 100 gallons of water.

Automatic proportioners - To make stock solution, add contents of 5 packets to 2 gallons of water for chickens and to 4 gallons of water for turkeys. Set proportioner to feed at rate of 1 fl oz of stock solution per gallon of water.

Treatment Period - 5 consecutive days.

Dairy Calves, Dairy Heifers, and Beef Cattle
Dosage: 25mg/lb first day followed by 12.5 mg/lb/day for 4 days.

Amount of Stock Solution for Cattle*	Sulfadimethoxine in Water Water Consumption	
	(Summer)	(Winter)
First Day Add	1 qt 10 gal	7 gal
	2 qt 20 gal	14 gal
	1 gal 40 gal	28 gal
Next 4 Days Add	1 qt 20 gal	14 gal
	2 qt 40 gal	28 gal
	1 gal 80 gal	56 gal

*Note: Make a cattle stock solution by adding 1 packet of Sulfadimethoxine Soluble Powder to 1 gal of water.
** This dosage recommendation is based on a water consumption of 1 gal per 100 lb of body weight per day, the expected water consumption rate for summer. Water consumption during cold months (winter) may drop markedly (50-60%). Accordingly, adjustments must be made in the dilution rates to compensate for this and insure proper drug intake.
For treatment of individual cattle, Sulfadimethoxine Soluble Powder stock solution for cattle may be given as a drench. Administer using same mg/lb dosage as outlined above.
*Twenty fluid ounces of cattle stock solution will medicate one-600 lb animal initially or twelve-600 lb animals on maintenance dose.
Contents of packet will medicate six-600 lb animals initially or twelve-600 lb animals on maintenance dose.

Treatment Period - 5 consecutive days

Caution: Chickens and Turkeys - If animals show no improvement within 5 days, discontinue treatment and re-evaluate diagnosis. Prepare a fresh stock solution daily. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication. Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders. Cattle - During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, re-evaluate diagnosis. Treatment should not be continued beyond 5 days.

Residue Warning:
Chickens and Turkeys - Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.
Cattle - Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves.
Do not use in calves to be processed for veal.

CONTACT INFORMATION: To report adverse drug events, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), please contact First Priority, Inc. at (800) 650-4899 or www.prioritycare.com.
For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/reportanimalad>.

STORAGE: Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.

Lot No. Exp. Date
NO VARNISH

Priority Care is a registered trademark of First Priority, Inc.

For a complete list of Priority Care distributors, please call our Customer Service Department at:
1-800-650-4899, or visit our Website: www.prioritycare.com